

Full Paper

## Copper(II) Complexes with Ligands Derived from 4-Amino-2,3-dimethyl-1-phenyl-3-pyrazolin-5-one: Synthesis and Biological Activity

Tudor Rosu <sup>1\*</sup>, Simona Pasculescu <sup>1</sup>, Veronica Lazar <sup>2</sup>, Carmen Chifiriuc <sup>2</sup> and Raluca Cernat <sup>2</sup>

<sup>1</sup> Faculty of Chemistry, Department of Inorganic Chemistry, University of Bucharest, 050107, Romania. Fax: (040) 21 3159249. E-mail: Simona Pasculescu: simonapasc@yahoo.com

<sup>2</sup> Faculty of Biology, Department of Microbiology, University of Bucharest, 050107, Romania. Tel. (040) 21 3181576. E-mails: Veronica Lazar: lazar@botanic.unibuc.ro; Carmen Balotescu: carmen\_balotescu@yahoo.com; Ramona Cernat: cernat@botanic.unibuc.ro

\* Author to whom correspondence should be addressed; E-mail: rosu.tudor@unibuc.ro or t\_rosu0101@yahoo.com

Received: 9 August 2006; in revised form: 22 September 2006 / Accepted: 5 November 2006 / Published: 17 November 2006

---

**Abstract:** The synthesis of Cu(II) complexes derived from Schiff base ligands obtained by the condensation of 2-hydroxybenzaldehyde or terephthalic aldehyde with 4-aminoantipyrine (4-amino-2,3-dimethyl-1-phenyl-3-pyrazolin-5-one) is presented. The newly prepared compounds were characterized by <sup>1</sup>H-NMR, UV-VIS, IR and ESR spectroscopy. The determination of the antimicrobial activity of the ligands and of the complexes was carried out on samples of *Escherichia coli*, *Klebsiella pneumoniae*, *Acinetobacter boumanii*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Candida sp.* The qualitative and quantitative antimicrobial activity test results proved that all the prepared complexes are very active, especially against samples of *Ps. aeruginosa*, *A. Boumanii*, *E. coli* and *S. aureus*.


**Keywords:** Schiff bases, antimicrobial activity, copper complexes, aminoantipyrine.

---

## Introduction

The well-known Cu(II) ion forms a series of coordination compounds with well defined structures. It plays an important role in the numerous biological processes that involve electron transfer reactions or the activation of some anti-tumor substances [1]. Copper is an essential micronutrient for feeding and a co-factor of several enzymes involved in oxidative metabolism:  $\beta$ -hydroxylases, quercetinase, ceruloplasmine, cytochromoxidase, mono-aminoxidase, superoxydismutase, ascorbic acid oxidase and tyrosinase. The catalytic role of these enzymes is the result of two processes: a) the reduction of the  $\text{Cu}^{2+}$  cation to  $\text{Cu}^+$ ; b) the fixation of the molecular oxygen [2]. As a cofactor of ceruloplasmin, copper contributes to the oxidation of Fe(II) to the corresponding Fe(III) form. Being related to transferin, the latter may cross the cell membranes [3]. Copper also has functions in erythropoiesis and hemoglobin-gensis, favoring, together with molybdenum, intestinal absorption, sediment mobilization and increases in plasmatic iron levels. Apart from its numerous functions in metabolic processes, copper is also recognized as a part in the immune function [3]. Superoxydismutase, which transforms toxic superoxide radicals in oxygen and peroxide, is dependent on copper and zinc. The  $\text{Cu}^{2+}$  ion is involved in the expression of genes for the metal-binding proteins [3] and it is also found in copper-protein combinations displaying a pseudotetrahedral symmetry and having effects in bio-systems. Through aminoxidase, copper interferes in the metabolism of the conjunctive tissue, contributing to the trophicity of vascular sides [4-8]. Taking into account the daily necessary quantity of Cu(II) in the organism (2-3 mg/day), its distribution and metabolism in the organism, toxicity, numerous simple or complex combinations of copper are used in the treatment of a variety of diseases, including inflammatory processes, cancer, ulcers, nervous system and heart diseases. This paper presents the synthesis and characterization of Cu(II) complexes with Schiff bases obtained by the condensation of 4-amino-2,3-dimethyl-1-phenyl-3-pyrazolin-5-one with 2-hydroxybenzaldehyde and terephthalic aldehyde, respectively (Figure 1).

**Figure 1.** Ligand structures.



Furthermore, and taking into consideration the use of copper complexes in the treatment of some diseases, mentioned above, we have tested the antimicrobial activity of the prepared ligands and complexes using strains of *Escherichia coli*, *Klebsiella pneumoniae*, *Acinetobacter boumanii*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Candida albicans* and *Candida tropicalis* isolated from different pathological products from patients with infections associated with the use of cardiovascular prosthetic devices. The antimicrobial activity of the Schiff bases of antipyrine and their complexes have been discussed previously [9-16].

## Results and Discussion

### Synthesis


The Schiff base ligands **ASAAP** and **ATAAP** were prepared by the condensation in methanol of one or two molecules of 4-amino-2,3-dimethyl-1-phenyl-3-pyrazolin-5-one with 2-hydroxybenzaldehyde and terephthalic aldehyde, respectively. The Cu complexes  $\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2\text{Cl}$  (**a**) and  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  (**b**) were then prepared by reaction of the appropriate ligands with aqueous solutions of a suitable Cu(II) salt.

### Properties

The molar conductivity value ( $32.7 \Omega^{-1}\cdot\text{cm}^2\cdot\text{mol}^{-1}$ ) of the  $\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2\text{Cl}$  complex in nitrobenzene indicates that this compound is a 1:1 electrolyte, while the molar conductivity value below  $10 \Omega^{-1}\cdot\text{cm}^2\cdot\text{mol}^{-1}$  in nitrobenzene for the  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  complex indicates that the latter is a non-electrolyte.

The complexes were also investigated by thermo-gravimetry (TG). Experimental data for these analyses are presented in Figure 2. The weight loss between 125–158°C for the  $\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2\text{Cl}$  complex is attributed to the loss of two water molecules per molecule of complex.

**Figure 2.** TG, DTG and ATD curves for:  $\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2\text{Cl}$  (**a**);  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  (**b**).



The second step (410–670°C) corresponds to the elimination of one molecule of ligand per molecule of complex. The thermal analysis curve of the  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  complex presents two weight loss steps (196–270°C and 380–694°C). The final residue was analyzed by IR spectroscopy and was identified as CuO and the % Cu corresponded to the calculated one.

### IR Spectra

Compared to the IR spectra of the ligands, it was observed that the frequency of the specific band of the  $\nu_{\text{C}=\text{N}}$  bond ( $1653, 1654 \text{ cm}^{-1}$ ) is moved towards lower wavenumbers by approx.  $13 - 15 \text{ cm}^{-1}$  in the spectra of the complexes, which confirms the coordination of the nitrogen atom to the metallic ion. In the IR spectrum of the **ASAAP** ligand, a wide medium intensity band occurs in the  $3210\text{--}3370 \text{ cm}^{-1}$

range, along with a narrow band of medium intensity ( $1140\text{ cm}^{-1}$ ), assigned to the phenolic  $\text{-OH}$  groups. In the IR spectrum of complex **(a)** the first band disappears and the second band moves towards lower wavenumbers, indicating coordination of the ligand to the  $\text{Cu}^{2+}$  ion through the phenolic  $\text{-OH}$  group oxygen. In this spectrum two narrow intense bands at  $897\text{ cm}^{-1}$  and  $572\text{ cm}^{-1}$  specific for coordinated water molecules [17] are also seen. For both complexes the specific  $\nu_{>\text{C}=\text{O}}$  band (cyclic keto group present in the pyrazolone ring:  $1615, 1596\text{ cm}^{-1}$ ) moves towards lower wavenumbers ( $1589, 1580\text{ cm}^{-1}$ ), suggesting the coordination of the ligand to the metallic ion via the  $>\text{C}=\text{O}$  group.

**Table 1:** IR spectral data for the prepared ligands and complexes.

Compound	$\nu_{\text{C}=\text{N}}$	$\nu_{\text{Ar-OH}}$	$\nu_{>\text{C}=\text{O}}$	$\rho_{\text{r}}$	$\rho_{\text{w}}$	$\nu_{\text{SO}_4^{2-}}$			
						$\nu_1$	$\nu_2$	$\nu_3$	$\nu_4$
$\text{C}_{18}\text{H}_{17}\text{N}_3\text{O}_2$	1653	3210- 3370 1140	1615	-	-	-	-	-	-
$[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$ <b>(a)</b>	1640	- 1120	1594	897	572	-	-	-	-
$\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2$	1654	-	1595		-	-	-	-	-
$[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$ <b>(b)</b>	1639	-	1565		-	995	462	1105 1170	574 610 641

Moreover, in the spectrum of complex **(b)** a characteristic band corresponding to the bidentate coordination of the  $\text{SO}_4^{2-}$  ion appears. Thus, the  $\nu_1$  and  $\nu_2$  frequencies specific to a  $T_d$  arrangement appear as medium intensity bands, and the  $\nu_3$  and  $\nu_4$  frequencies each split into three bands, which suggest a low symmetry, probably reduced towards  $\text{C}_{2v}$  [17].

#### Electronic spectra.

As seen from the data presented in Table 2, the  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$  complex presents a single absorption band at  $14600\text{ cm}^{-1}$  corresponding to the  $d-d$  transition, indicating the low  $\text{C}_{2v}$  symmetry of the  $\text{Cu}^{2+}$  ion [18,19]. The brown colored  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  complex presents two types of  $d-d$  type transitions, whose values are characteristic of a deformed tetrahedral symmetry, with the term of the fundamental state  $d_{xy}$  [19].


**Table 2.** Electronic spectra of the synthesized complex combinations.

Compound	Transitions $d-d$ ( $\text{cm}^{-1}$ )		Geometry
$[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$	$x^2-y^2 \rightarrow xz; yz; xy$ 14600		$\text{C}_{2v}$
$[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$	-	$xy \rightarrow xz; yz$ 13570	$\text{Td}$ deformed

## ESR spectra


The ESR spectral data concerning the  $\text{Cu}^{2+}$  ion in tetrahedral symmetry are relatively poor. These present a special interest due to the fact that, in the case of hyperfine interaction, the values of the  $A$  constants are considerably lower than the observed ones for a complex of  $\text{Cu}(\text{II})$  with a  $\text{O}_h$  or  $\text{D}_{4h}$  geometry, and the values of the  $\{g\}$  tensor are higher [20-24]. The room temperature ESR spectrum of the complex  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})]\text{Cl}$ , as a powder, is presented in Figure 3.

**Figure 3.** ESR for  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$ , as a powder, at room temperature.



The values of the  $\{g\}$  tensor confirm the  $\text{C}_{2v}$  symmetry of the  $\text{Cu}^{2+}$  pentacoordinated ion [24]. For the  $\text{Cu}(\text{II})$  complex with low symmetry, the fundamental state for the paramagnetic electron is not described by a single  $d$  function, but there is a mixture of them. The mixture degree of these functions increases as the symmetry decreases [24]. The high values of the  $\{g\}$  tensor confirm a low symmetry. The ESR spectrum for the complex  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  as a powder, at room temperature, is presented in Figure 4.


**Figure 4.** ESR for  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{S}_2\text{O}_{10})]$ , as a powder, at room temperature.



The  $\{g\}$  tensor values,  $g_1 = 2.088$  and  $g_2 = 2.406$ , confirm the  $\text{T}_d$  symmetry [20-23]. As noted, the spectrum does not present interactions that would indicate a hyperfine structure. The lack of hyperfine structure or the presence of very small hyperfine splitting for the  $\text{Cu}^{2+}$  ion in  $\text{T}_d$  or lower symmetry were explained by C.A. Bates *et al.*, by mixing the  $4p_z$  orbital of the metallic ion with the  $3d_{xy}$  orbital which defines the fundamental state [20]. The values of the electronic transitions, of the  $\{g\}$  tensor,

and IR spectral data lead to the conclusion that the  $\text{Cu}^{2+}$  ion of the  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$  complex is pentacoordinated with a  $\text{C}_{2v}$  symmetry (Figure 5a) [24], while in the  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  complex the  $\text{Cu}^{2+}$  ion has a deformed tetrahedral geometry (Figure 5b) [19].


**Figure 5.** Structures of the complexes  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$  (a) and  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  (b).



#### Antimicrobial activity assays


The antimicrobial activity of the complexes and ligands was screened by adapted qualitative, diffusimetric methods (i.e. distribution of the tested solutions on filter paper disks, in agar wells or in spots on solid media that have been inoculated with test microbial strains) and quantitative methods based on serial two-fold dilutions of the tested compounds in order to establish the corresponding Minimal Inhibitory Concentrations (MIC). Five bacterial strains, i.e. *Escherichia coli*, *Klebsiella pneumoniae*, *Acinetobacter boumanii*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and two fungal strains, i.e. *Candida albicans* and *Candida tropicalis*, freshly isolated from different clinical sources from patients with infections associated with the use of cardiovascular prosthetic devices and identified by conventional methods were cultivated on solid media and incubated at  $37^\circ\text{C}$  for 24 hrs prior to testing.

**Figure 6.** Appearance of the qualitative screening test showing the activity of  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$  by the three adapted agar diffusion methods.



The qualitative screening results demonstrated that the three examined diffusion methods all exhibited different sensitivities in detecting the antimicrobial potential of the tested compounds. The most efficient one for the different bacterial strains proved to be the spot method, as exemplified in Figure 6. The quantitative assay results (Figure 7) showed that the tested compounds exhibited variable MICs and selective antimicrobial activity, depending on the microbial strains. All tested compounds proved to be active on *Ps. aeruginosa*, well known for its high constitutive and acquired resistance rates. The Schiff base **ASAAP** and the complex  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$  exhibited high bactericidal activity towards *E. coli* and *A. Boumanii*, while the Schiff base ligands **ASAAP** and **ATAAP** and the  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  complex also showed good activity against *S. aureus*, *Ps. aeruginosa* and *E. coli*, proving their potential usefulness as broad spectrum antimicrobial agents.

**Figure 7.** The graphic representation of the MIC values (mg/mL) of the tested compounds towards different bacterial strains.



## Conclusions

The IR, electronic transition and  $\{g\}$  tensor value data lead to the conclusion that the  $\text{Cu}^{2+}$  ion in the complex  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$  is pentacoordinated with a  $\text{C}_{2v}$  symmetry (Figure 5a), whereas in the complex  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$  the  $\text{Cu}^{2+}$  ion has a deformed tetrahedral geometry (Figure 5b). The sensitivity spectrum of the microbial strains towards the ligands and the corresponding complexes was determined by qualitative and quantitative methods and the following conclusions were reached:

- the qualitative anti-microbial activity screening results of the tested compounds proved that the most efficient test method was the spot method.
- the quantitative anti-microbial activity test results proved that both the ligands and the complex combinations have specific anti-microbial activity, depending on the microbial species tested.

## Experimental

### General

The reagents used in this work were commercial products (Merck and Chimopar Bucuresti). Electronic spectra were recorded using a Jasco V-550 spectrophotometer, in diffuse reflectance mode, using MgO dilution matrices. IR spectra (KBr pellets) were recorded in the 4000-400  $\text{cm}^{-1}$  region with a BioRad FTS 135 spectrophotometer. ESR spectra were recorded on a ART-6 model IFA-Bucuresti type spectrophotometer, equipped with a field modulation unit at 100kHz. The measurements were done in the X band, on micro-crystalline powder at room temperature using DPPH as standard. The  $^1\text{H-NMR}$  spectra were recorded using a Bruker DRX 400 spectrometer. Chemical elemental analyses were done with a Carlo-Erba LA-118 microdosimeter (for C, N) and an AAS-1N Carl-Zeiss-Jena spectrometer [Cu(II)], respectively. Chlorine was determined by gravimetric analysis. The complexes were studied by thermo-gravimetry (TG) in a static nitrogen atmosphere, with a sample heating rate of  $10^\circ\text{C}/\text{min.}$ , using a DuPont 2000 ATG thermobalance. Molar conductances of the complexes were measured in nitrobenzene at room temperature using a Consort type C-533 conductivity instrument.

### Synthesis of the Schiff base 1-phenyl-2,3-dimethyl-4-(*N*-salicylidene)-3-pyrazolin-5-one (ASAAP)

A solution of 4-amino-2,3-dimethyl-1-phenyl-3-pyrazolin-5-one (0.203 g, 1 mmol) in methanol (20 mL) was added to a solution of *o*-hydroxybenzaldehyde (0.15 mL, 1 mmol) in methanol (10 mL). The mixture obtained was refluxed for an hour, then stirred for 3 hrs at room temperature and left at the same temperature for a day. The resulting intense yellow colored precipitate was filtered, washed with methanol and dried. Elemental analysis: Calc. C%, 70.35; N%, 13.66; Found. C%, 71.16; N%, 13.21. IR:  $1653\text{ cm}^{-1}$  (azomethine group) [8,9];  $^1\text{H-NMR}$  spectra [ $\text{CDCl}_3$ ,  $\delta$  (ppm),  $J$  (Hz)]:  $\delta = 2.41$  (s, 3H, - $\text{CH}_3$ ); 3.16 (s, 3H, - $\text{CH}_3$ ); 6.90 (m, 1H, H-Ar); 6.95 (d, 1H, H-Ar); 7.24-7.56 (m, 7H, H-Ar); 9.84 (s, H, - $\text{N}=\text{CH}$ -);  $\delta > 7$  (H-OH).

### Synthesis of the Schiff base bis(1-phenyl-2,3-dimethyl-3-pyrazolin-5-one-4-imino) terephthalic aldehyde (ATAAP)

A solution of 4-amino-2,3-dimethyl-1-phenyl-3-pyrazolin-5-one (0.253 g, 1.25 mmol) in methanol (15 mL) was added to a solution of terephthalic aldehyde (0.067 g, 0.5 mmol) in methanol (15 mL). The resulting yellow solution was refluxed for 30 minutes and then left at room temperature for approx. 6 hrs. The intense yellow precipitate formed was filtered, washed with methanol and dried. Elemental analysis: Calc. C%, 71.42; N%, 16.66; Found. C%, 72.11; N%, 16.21; IR:  $1654\text{ cm}^{-1}$  (azomethine group) [8,9].  $^1\text{H-NMR}$  spectra [ $\text{CDCl}_3$ ,  $\delta$  (ppm),  $J$  (Hz)]:  $\delta = 9.61$  (s, 2H, - $\text{N}=\text{CH}$ -); 7.88 (s, 4H, Ar-H 1,4- disubstituted); 7.57-7.37 (m, 10H, Ar-H); 3.21 (s, 6H, N- $\text{CH}_3$ );  $\approx 2.5$  (s, 6H, - $\text{CH}_3$ ).

### Synthesis of the complex $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$ (a)

A methanol solution (15 mL) of ASAAP (0.307 g, 1 mmol) was added to  $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$  (0.170 g, 1 mmol) dissolved in distilled water (10 mL). This solution was refluxed for 2 hrs and left at room

temperature for three days. A brown-red precipitate was formed, which was filtered, washed with ethanol and dried. The elemental analysis results (calc.: C%, 48.81; N%, 9.49; Cu%, 14.46; Cl%, 8.04; found: C%, 49.33; N%, 9.09; Cu%, 14.25; Cl%, 7.83) confirm the molecular formula  $[\text{Cu}(\text{C}_{18}\text{H}_{16}\text{N}_3\text{O}_2)(\text{H}_2\text{O})_2]\text{Cl}$ .

#### *Synthesis of the complex $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$ (b)*

A DMF solution (15 mL) of **ATAAP** (0.252g, 0.5 mmol) was added to  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  (0.250 g, 1 mmol) dissolved in distilled water (15 mL). The resulting green solution was refluxed for two hours, during which time the green color turned brown. This solution was left at room temperature for four days. A brown precipitate was formed, which was filtered, washed with ethanol and dried. The elemental analysis data (calc.: C%, 43.69; N%, 10.19; Cu%, 15.53; found: C% 44.12; N%, 16.11; Cu%, 15.24) confirms the molecular formula  $[\text{Cu}_2(\text{C}_{30}\text{H}_{28}\text{N}_6\text{O}_2)(\text{SO}_4)_2]$ .

#### *Biological assays*

The fresh cultures obtained from clinical isolates were suspended in sterile saline and adjusted to a standard density of 0.5 MacFarland. The microbial suspensions were plated on solid Mueller Hinton medium and solutions of the test compounds (10  $\mu\text{L}$ ) prepared in DMF (1 mg/mL) were added on filter paper disks, in agar wells or in spots. Concomitantly, the disks were impregnated with the same concentration of gentamycin, which was used as reference standard for reporting the antibiotic sensitivity. The plates were incubated at 37°C for 24 hrs. During incubation, the tested compounds diffused around the test area creating a concentration gradient. The antimicrobial activity was recorded as any area of microbial growth inhibition that occurred in the diffusion area. The quantitative antimicrobial activity assays were performed by the two-fold serial microdilution method in liquid medium (nutrient broth for bacterial and liquid YPG for fungal strains). Serial two-fold dilutions of a stock solution of test compound in DMF (from 1000 to 62.5  $\mu\text{g}/\text{mL}$ ) were performed in 60 multi-well plates, in a total volume of 200  $\mu\text{L}$  medium and standard microbial suspension (50  $\mu\text{L}$ ) was added in each well. After 18-24 hours, the plates are examined visually for evidence of bacterial growth. Results are recorded as minimum inhibitory concentrations (MIC) at the highest dilution (lowest concentration) of the tested compound that completely inhibited microbial growth.

#### **Acknowledgements**

The authors thank Organic Chemistry Department - ICECHIM, Bucharest for microanalysis and IFIN - Horia Hulubei, Bucharest, Romania, for help with EPR spectroscopy.

#### **References**

1. Karlin, K.D.; Gultneh, Y. Bioinorganic chemical modeling of dioxygen-activating copper proteins. *J. Chem. Ed.* **1985**, *62*, 983-992.

2. Halfen, J.A.; Mahapatra, S.; Wilkinson, E. C.; Kaderli, S.; Young, Jr. V. G.; Que, Jr. L.; Zuberbühler, A.D.; Tolman W.B. Reversible cleavage and formation of dioxygen O-O bond within a copper complex. *Science* **1996**, *271*, 1397-1400.
3. a) Berdanier, C.D.; Groff J.L.; Gropper, S.S. *Advanced Nutrition and Human Metabolism*, 3<sup>rd</sup> Ed.; Wadsworth/Thomson Learning: Belmont, CA, **1999**; pp. 87-105; b) Gropper, S.S.; Smith, J.L.; Groff, J.L. *Advanced Nutrition and Human Metabolism*, 4<sup>th</sup> Ed.; Thomson Wadsworth Publishing Co.: Belmont, CA, **2005**.
4. Brill, A.S.; Martin, R.B.; Williams R.J.P. *Electronic Aspects of Biochemistry*; Academic Press Inc.: New-York, **1964**; pp. 519–557.
5. Frieden, E.; Osaki, S.; Kobayashi H. Copper proteins and oxygen. Correlations between structure and function of the copper oxidases. *J. Gen. Physiol.* **1965**, *49 (1 Suppl.)*, 213-252.
6. Sugiura, Y.; Hirayama, Y.; Tanaka, H.; Ishizu, K. Copper(II) complex of sulfur-containing peptides. Characterization and similarity of electron spin resonance spectrum to the chromophore in blue copper proteins. *J. Am. Chem. Soc.* **1975**, *97*, 5577-5581.
7. Fee, J.A. Copper proteins - Systems containing the "blue" copper center. *Structure and Bonding*; Springer-Verlag: Heidelberg, **1975**; Vol. 23, pp. 1-60.
8. Sakaguchi U.; Addison, A.W. Structural implications for blue protein copper centers from electron spin resonance spectra of copper sulfide (Cu<sup>II</sup>S<sub>4</sub>) chromophores. *J. Am. Chem. Soc.* **1977**, *99*, 5189–5190.
9. Raman, N., Kulandaisamy, A.; Shunmugasundaram, A.; Jeyasubramanian, K. Synthesis, spectral, redox and antimicrobial activities of Schiff base complexes derived from 1-phenyl-2,3-dimethyl-4-aminopyrazol-5-one and acetoacetanilide. *Transit. Metal Chem.* **2001**, *26*, 131-135.
10. Raman, N.; Kulandaisamy, A.; Jeyasubramanian, K. Synthesis, spectral, redox and antimicrobial activity of Schiff base transition metal(II) complexes derived from 4-aminoantipyrine and benzil. *Synth. React. Inorg. Met-Org. Nano-Met. Chem.* **2002**, *32*, 1583-1610.
11. Singh, L.; Sharma, D.K.; Singh, U.; Kumar, A. Synthesis and spectral studies of Cu(II) coordination compounds of 4[N-(cinnamalidene)amino]antipyrine semicarbazone. *Asian J. Chem.* **2004**, *16*, 577-580.
12. Raman, N.; Thangaraja, C.; Johnsonraja, S. Synthesis, spectral characterization, redox and antimicrobial activity of Schiff base transition metal(II) complexes derived from 4-aminoantipyrine and 3-salicylideneacetylacetone. *Centr. Eur. J. Chem.* **2005**, *3*, 537-555.
13. Pandey, O.P.; Sengupta, S.K.; Dwivedi, A. Organophosphorus derivatives containing antipyrine ring as chemotherapeutics against fungal pathogens of sugarcane. *Electron. J. Environ. Agr. Food Chem.* **2005**, *4*, 886-891.
14. Agarwal, R.K.; Singh, L.; Sharma, D. K. Synthesis, spectral and biological properties of copper (II) complexes of thiosemicarbazones of Schiff bases derived from 4-aminoantipyrine and aromatic aldehydes. *Bioinorg. Chem. Appl.* **2006**, article ID 59509.
15. Agarwal, R.K.; Gargb, R.K.; Sindhub, S.K. Synthesis and magneto-spectral investigations of some six and nine coordinated complexes of lanthanides(III) derived from 4[N-(2'-hydroxy-1'-naphthalidene)amino]antipyrine thiosemicarbazone. *J. Iran. Chem. Soc.* **2005**, *2*, 203-211.
16. Agarwal R. K.; Prasad S. Synthesis and spectral investigations of some platinum metals ions coordination compounds of 4[N-(furan-2'-carboxalidene)amino]antipyrine thiosemicarbazone

- and 4[N-(3',4',5'-trimethoxybenzalidene)amino]antipyrine thiosemicarbazone. *Turk. J. Chem.* **2005**, *29*, 289-29.
17. Nakamoto K. *Infrared Spectra of Inorganic and Coordination Compounds*; Wiley and Sons: New York, **1986**; pp. 248-249.
  18. Tyagi, S.; Hathaway, B.J. Crystal structure and electronic properties of bis(2,2'-bipyridyl)-cyanocopper(II) nitrate dihydrate: a correlation of the in-plane angular distortion with the splitting of the electronic spectrum. *J. Chem. Soc. Dalton Trans.* **1983**, 199-203.
  19. Lever, A.B.P. *Inorganic Electronic Spectroscopy*, 2<sup>nd</sup> Edn; Elsevier Science: New York, **1984**; pp. 560-571.
  20. Bates, C.A.; Smoore, W.; Standley, K.J.; Stevens, K.W.H. Paramagnetic resonance of a Cu<sup>2+</sup> ion in a tetrahedral crystal field. *Proc. Phys. Soc.* **1962**, *79*, 73-83.
  21. Sharnoff, M. Electron paramagnetic resonance and the primarily 3d wavefunctions of the tetrachlorocuprate ion. *J. Chem. Phys.* **1965**, *42*, 3383-3395.
  22. Forster, D.; Weiss, V.W. An electron paramagnetic resonance study of cupric ion tetrahedrally coordinated by nitrogen atoms. *J. Phys. Chem.* **1968**, *72*, 2669-2671.
  23. Yokoi, H.; Addison, A.W. Spectroscopic and redox properties of pseudotetrahedral copper(II) complexes. Their relation to copper proteins. *Inorg. Chem.* **1977**, *16*, 1341-1349.
  24. Wasson, J.U.R.; Klassen, D.M.; Richardson, H.W.; Hatfield, W.E. Low-symmetry copper(II) complexes. Spectral properties of dihalo[2,6-di(2'-quinolyl)pyridine]copper(II) complexes. *Inorg. Chem.* **1977**, *16*, 1906-1911.

*Sample availability:* Samples of the ligands **ASAAP** and **ATAAP** and the complexes **(a)** and **(b)** are available from MDPI.