

The Effects of Conjugation Path Variation on Electron Delocalization in Phenoxy-Based Systems.

Yanbing Liu and Paul M. Lahti*

Department of Chemistry, University of Massachusetts, Amherst, MA 01003 USA. Tel.: (+1) 413-545-4890, Fax: (+1) 413-545-4490

* Author to whom correspondence should be addressed; e-mail: lahti@chem.umass.edu.

Received: 23 April 2004 / Accepted: 2 June 2004 / Published: 31 August 2004

Abstract: A number of persistent 2,6-di-*tert*-butylphenoxy based radicals were synthesized as models for unpaired spin delocalization as a function of conjugation pathway, and were investigated by ESR and UV-vis spectroscopy. Phenoxy spin delocalizes significantly onto a *para*-phenyl ring, but further delocalization through a *meta*-vinyl or *meta*-carbonyl linkage is not detectable by ESR hyperfine coupling. UV-vis spectra do show a red shift of the longest wavelength transition for a carbonyl-type substituent by comparison to a vinyl substituent in the *meta*-position of a *para*-phenyl group. By comparison, a *para*-styryl substituent on the phenoxy causes very large spin delocalization from the phenoxy unit, with the largest hyperfine coupling being found on the ethenyl unit of the assemblage. This large delocalization accounts for the reactivity of radicals that incorporate such units.

Keywords: 2,6-Di-*tert*-butylphenoxy, conjugated polyradicals, spin delocalization, exchange, molecular magnetism, electron spin resonance, hyperfine coupling.

Introduction


Recent interest in molecule-based magnetic materials has spurred much work upon organic diradical and polyradical systems [1]. Rajca's review of this area subdivided organic polyradicals into macrocyclic and pendant spin-bearing systems [2]. The former category of systems has been elegantly exploited by Rajca's group to make very high spin polyradicals with total spin quantum numbers (S) that surpass those of inorganic elements, and have even been shown to order [3] as hydrocarbon-based materials. Pendant systems have simultaneously been much more studied, but disappointing until recently [4]. Computational studies have shown that pendant systems have an inherently smaller intramolecular exchange than the macrocyclic backbone conjugated systems, such that the exchange interaction in the pendants is easily decreased or virtually eliminated by torsional deconjugation of the pi-network [5]. However, if planarity can be sufficiently maintained to allow significant intramolecular exchange, pendant systems are tolerant of some spin defects because the pathways of conjugation between unpaired spins are not blocked [6]. Recently, Nishide's group has synthesized star polyradicals with two-dimensional architecture [7] somewhat similar in concept to the related Rajca-type polyarylmethyl backbone polyradicals and obtained magnetic force microscopy measurements [4f-g] for these systems. Such approaches reinforce the notion that variation and control of structure can give desirable and extended exchange behavior in purely organic systems.

We have for some time investigated the incorporation of phenoxy radicals into polyradical systems generated by photochemical and solution oxidation methods [8]. In this study, we describe the synthesis and electron spin resonance properties of a set of phenoxy-based models to investigate some of the structure-property relationships that affect and enhance spin delocalization and exchange in polyradicals that incorporate phenoxy spin-carrying units.

Results and Discussion


Our studies focused on the extent of delocalization for the phenoxy group's unpaired spin density in systems with *para*-phenyl, *para*-vinylene and related substituents. Nishide's success [4e-f] in showing magnetic exchange for polyphenylenevinylenes with pendant phenoxy radicals impelled our desire to determine how well delocalization occurs in such systems. Electron spin resonance hyperfine coupling (ESR hfc) results from such polyradicals give some information at low degrees of oxidization from precursor polyphenols to polyradicals, but the polyradicals themselves tend to have broad, featureless ESR spectra. We therefore investigated the ESR spectra for two main model systems and a set of building blocks obtained during synthesis of the main models.

Model system **1** was pursued to imitate putative polyradical **2**, an all *meta*-linked system. System **2** should have at best a small preference for a high spin state due to its connectivity [1]. Although each monomer unit in **2** has an unpaired electron, with connectivity such that the number of unpaired spins nominally is equal to the degree of polymerization ($n^* > n^\circ$ in Figure 1)[1b], **2** can be considered disjoint [1c] due to the localization of molecular spin orbitals (connections only through nodal sites).



Despite their low exchange energies between unpaired electrons, such near-degenerate open-shell systems have recently become interesting due to the possibilities for spintronic [9] strategies to “tune” their magnetic nature by application of electrical fields. We aimed to determine the stability of **1**, and to find out whether delocalization was sufficient to produce an ESR-observable triplet diradical state, indicating significant exchange.


Figure 1. Parity based analysis for polymers **2** and **4**. Upper two structures show polymer **2**, parsed into segments by comparison model system **1** and model systems in Table 1 below. Lower structure shows polymer **4** parsed into stilbenic units related to **5**.



Model system **3** was pursued to model polyradical **4**, in which a phenoxy system would be directly incorporated into a polyphenylenevinylene chain. System **4** is expected to exhibit strong delocalization due to nondisjoint connectivity. We aimed to test its spin delocalization and stability. The related stilbenoxy system **5** has been known for some time [10], and constitutes a useful comparison to **3**, as we shall see below.

As part of investigating exchange in **1-2**, we made a number of model phenols fragments with more limited conjugation, as shown in Figure 2. 3,5-Di-*tert*-butyltrimethylsiloxyphenyl boronic acid (**6**) was coupled with 3-iodobenzaldehyde by the Suzuki method to give **7** (under our reaction conditions, Suzuki couplings with **6** always gave products in which the trimethylsilyl group was cleaved). Compound **7** in turn was converted to vinyl analog **8** by a Wittig reaction. Compound **6** was also coupled with 5-iodoisophthalaldehyde by the Suzuki method, to give dialdehyde **9**, which was subjected to double Wittig vinylation to yield **10**. The analogous couplings with 5-iodoisophthalic acid and its dimethyl ester yielded phenols **11** and **12**. These compounds constitute a set of limited-conjugation phenoxy precursors, all having *para*-phenyl substituents with *meta*-linkage extension.

Figure 2. Synthesis of fragment model segments related to polymer **2**.




Conditions: (a) K_2CO_3 , catalytic $Pd(OAc)_2/P(o\text{-tolyl})_3$, DMF, heat, 35%; (b) CH_3PPh_3Br , *tert*-Bu-Li, THF, 80%; (c) K_2CO_3 , catalytic $Pd(OAc)_2/P(o\text{-tolyl})_3$, DMF, heat, 42%; (d) CH_3PPh_3Br , *tert*-Bu-Li, THF, 81%; (e) K_2CO_3 , catalytic $Pd(OAc)_2/P(o\text{-tolyl})_3$, acetone-water, heat, 50%; (f) K_2CO_3 , catalytic $Pd(OAc)_2/P(o\text{-tolyl})_3$, DMF, heat, 77%.

Figure 3 shows the synthesis of the full structure precursor to **1**. 4-Bromo-2,5-di-*tert*-butylphenyl acetate was reacted with 4-methoxyphenyl boronic acid using palladium catalysis (Suzuki method) to give **13**, which was brominated to give **14** (even excess bromine only gave monobromination). Heck coupling of **14** with *meta*-divinylbenzene gave **15** in 70% yield. Lithium aluminum hydride treatment of **15** yielded diphenol **16**, the structure of which was confirmed by 1H -NMR and IR spectra. In


particular, its IR spectrum showed a strong and sharp absorption for the sterically hindered phenol OH at 3600 cm^{-1} , and an out-of-plane *trans*-HC=CH bend at 960 cm^{-1} . High-resolution mass spectrometry gave a molecular ion with $m/z = 750.4651$, in excellent agreement with the calculated value of 750.4648 for $\text{C}_{52}\text{H}_{62}\text{O}_4$.

Figure 3. Synthesis of precursor **16** to model system **1**.



Conditions: (a) K_2CO_3 , catalytic $\text{Pd}(\text{OAc})_2/\text{P}(\text{o-tolyl})_3$, DMF, heat, 64%; (b) Br_2 , CH_2Cl_2 , heat, 57%; (c) Bu_3N , catalytic $\text{Pd}(\text{OAc})_2/\text{P}(\text{o-tolyl})_3$, 1,3-divinylbenzene, DMF, heat, 44%; (d) LiAlH_4 , THF, 53%.

Figure 4. Synthesis of precursor **19** to model system **3**.




Conditions: (a) $\text{CH}_3\text{PPh}_3\text{Br}$, *n*-Bu-Li, THF, 60%; (b) Et_3N , catalytic $\text{Pd}(\text{OAc})_2/\text{P}(\text{o-tolyl})_3$, 1,3-diiodobenzene, DMF, heat, 70%; (c) LiAlH_4 , THF, 74%.

Figure 4 shows the synthesis of the diphenolic precursor to model system **3**. 3,5-Di-*tert*-butyl-4-hydroxybenzaldehyde was vinylylated by a Wittig reaction and acetyl-protected in a one-pot reaction to give **17**, which was subjected to Heck coupling with 1,3-diiodobenzene to give **18**. Subsequent deprotection yielded the diphenol **19**, the structure of which was confirmed by spectroscopic and elemental analysis.

Solution phase oxidation of diphenol **16** under argon with lead dioxide in benzene or toluene immediately gave a very deep blue solution with new UV-vis bands at 319, 372, and 614 (broad) nm and the solution ESR spectrum shown in Figure 5. The oxidation was carried out sufficient long to oxidize both phenol units in **16**, based on thin layer chromatographic analysis of the consumption of di-*tert*-butylphenol precursors in related oxidations (5-15 min). The spectrum was persistent for hours


under inert atmosphere or *in vacuo*. Analysis of the hyperfine coupling constants (hfc) was carried out using Duling's WINSIM program [11], showing major hfc on the phenoxyl ring and the *para*-connected phenyl ring as expected for alternant delocalization (see Figure 1 and Scheme 1).

Figure 5. Room temperature ESR spectrum derived from oxidation of **16** in toluene at 9.786 GHz



We observed no hfc indicating delocalization onto the *meta* vinyl unit or beyond, despite various attempts with different solution concentrations and at various temperatures from 200-300 K. We also were unable to resolve hfc from the C-H bond *ortho* to the methoxy group in the second phenyl ring, which should have a small negative spin density based on the spin distributions expected in Figure 1. This may be due to biradical-induced broadening that is unavoidable under our conditions, or possibly to overlap of multiple small hfc that are not resolved.



Scheme 1. Hfc assignments from Figure 5 for spectrum assigned to **1**, derived by solution oxidation of **16**; hfc given in gauss.



If biradical **1** possessed an exchange constant $J > a(H)$ – where $a(H)$ are the hfc of the aromatic protons – then we would expect its line-to-line spacing to be about 50% of that observed for analogous monophenoxyl radicals [12]. This was not observed, which would be consistent with $J < a(H)$. A toluene solution of **1** was frozen to 77 K to see whether evidence of a discrete biradical triplet state could be detected. No fine structure attributable to $\Delta M_s = 1$ transitions of a triplet state was observed

in the $g \sim 2$ region, but only a single peak with linewidth of $\Delta H_{pp} = 10\text{-}11$ gauss. No $\Delta M_s = 2$ transition was observed at half-field (1600-1700 gauss region of X-band spectrum) under any conditions attempted. These observations would be consistent with a triplet state having very small zero field splitting (zfs), but if so, any exchange between the radical units of **1** must be very small. As part of the investigations of systems related to polyradical **2**, we also obtained the ESR and UV-vis spectra derived by oxidation of the limited-conjugation systems with *meta*-linked extension, phenols **7-12** in Figure 2. These were tested both as potential building blocks for future studies, and because they possess nominally conjugating substitution in the same position as the repeat units for polyradical **2**. Table 1 summarizes the UV-vis and ESR spectral data derived from brief oxidation of these precursors to radicals. In this Table $a(H)$ are proton hyperfine constants in gauss, derived from on lineshape fitting using WINSIM [11].

Table 1: ESR hyperfine and UV-vis data for model phenoxyl radicals derived from phenols in Figure 2. All spectra obtained in benzene or toluene solutions at room temperature.


Compound	$a(H)/\text{gauss}$	UV-vis (λ_{max} , nm)
 $R = \text{CHO}$	$a_3 = 1.80$ (2 H) $a_5 = 0.68$ $a_4 = 1.60$ (2 H) $a_6 = 1.86$	350, 482, 599 (broad) dark green
$R = \text{CH}=\text{CH}_2$	$a_3 = 1.76$ (2 H) $a_5 = 0.69$ $a_4 = 1.70$ (2 H) $a_6 = 1.73$	328, 355, 507 red-purple
 $R = \text{CHO}$	$a_3 = 1.77$ (2 H) $a_6 = 1.78$ $a_4 = 1.65$ (2 H)	341, 458, 615(broad) dark green
$R = \text{CH}=\text{CH}_2$	$a_3 = 1.68$ (2 H) $a_6 = 1.92$ $a_4 = 1.64$ (2 H)	276, 337, 514 (broad) red-purple
$R = \text{CO}_2\text{H}$	$a_3 = 1.76$ (2 H)* $a_6 = 1.86$ $a_4 = 1.67$ (2 H)	267, 359, 505 (broad)* (yellow)
$R = \text{CO}_2\text{Me}$	$a_3 = 1.69$ (2 H) $a_6 = 1.70$ $a_4 = 1.69$ (2 H)	347, 475, 615 (broad) (blue)

*Very sparingly soluble in benzene, toluene

All radicals exhibited highly colored solutions that were persistent for hours if exposure to air was limited. All show significant hfc in both rings, showing delocalization from the phenoxyl onto the phenyl ring – however, in no case did we obtain resolvable hfc from the linking substituent in the *meta*-position (R in the table). The sum of all available information for both **1** and the radicals derived from **7-12** shows no more than 2-3% spin delocalization onto any π -orbital bearing carbon at the *meta*-position (R = vinyl, carbonyl carbons in Table 1), if our resolution is at least a conservative 0.5 G, and the McConnell relationship[13] holds such that $a(\text{H}) = (-22 \text{ G}) \times \rho(\pi\text{C})$, where $\rho(\pi\text{C}) = \pi$ -spin density of carbon to which an aryl C-H is attached with an hfc of $a(\text{H})$.

Although the ESR evidence shows neither significant delocalization of unpaired spin onto the *meta*-linked positions of **7-12**, nor major variation with substituent R (Table 1), the UV-vis spectra show a qualitative trend. The vinyl-substituted phenoxyl radicals derived from **8** and **10** show substantially blue-shifted long-wavelength bands relative to the carbonyl-substituted variants. The diacid system (R=CO₂H, Table 1) is yellow in benzene or toluene, but greenish in DMSO, so this phenoxyl may not fit the trend well due to its known hydrogen bonding in nonpolar solvents [14]. The trend may be due to differing charge-transfer character of the long wavelength band in the systems with the more electron-poor ring, namely those with the carbonyl substituents. In an effort to test a nondisjoint system where greater delocalization is expected, diphenol **19** was stirred with PbO₂ in benzene for 1-5 min to give deep yellow solutions. The color change was accompanied by appearance of a strong band in the UV-vis spectrum at 309 nm and quite weak bands at 457 and 488 nm. Oxidation of the precursor phenol groups was confirmed by disappearance of the OH stretching absorption at 3600 cm⁻¹ in the IR spectrum of a film of oxidized product. The ESR spectrum exhibited the multi-line pattern shown in Figure 7, which shows an excellent fit (correlation coeff = 0.993) to the following hfc: $a(\text{H}) = 6.37 \text{ G}$ (1 H), 2.93 (1 H), 1.69-1.79 (2 H), 1.40-1.48 (3 H), 0.53 G (1 H). The spectrum did not vary with the length of oxidation over 1-5 min, and persists for hours in the absence of air.


Figure 7. Room temperature ESR spectrum derived from oxidation of **19**. Experimental spectrum in toluene at 9.751 GHz (black); simulation (red) with WINSIM [11] using $a(\text{H}) = 6.37 \text{ G}$ (1 H), 2.93 (1 H), 1.69 (1 H), 1.59 (1 H), 1.48 (1 H), 1.47 (1 H), 1.40 (1 H), 0.53 G (1 H).



The spectrum from **19** exhibits an unusually large hfc of 6.4 G for one proton, requiring that the proton be attached at a position with a large spin population (nearly 30% based on the McConnell model described above). The hfc of the *meta*-hydrogen atoms on a *tert*-butylated phenoxy ring are typically 1-3 G. We therefore assigned the large hfc to a stilbenic olefin CH group, due to resonance structure **20A** (R=*tert*-Bu, Scheme 2). This is consistent with computations for 4-stilbenoxy model **20** (R=H) at the UB3LYP/EPR-II // UB3LYP/6-31G* level using Gaussian 98 [15]. The computations predict hyperfine couplings in good accord with those recently [16] reported for radical **20** (R=*tert*-Bu); the report gives one proton hfc of 6.39 G. Clearly, **20** and related systems are so delocalized that their major spin density site is not even found in the phenoxy ring, but instead on the olefinic group. This tendency is strengthened by the significant C=O bonding character in resonance structure **20A**.

The 4-stilbenoxyl **20** (R=*tert*-Bu) dimerizes readily in the solid state [10], although it shows a persistent solution ESR spectrum. We presume that the analogous system **3** is similarly reactive to form **21** or related species, explaining our inability to observe a triplet state spectrum for **3** despite expected strong exchange in this nondisjoint connectivity.


Scheme 2. Resonance structures of 4-stilbenoxyl radicals, and a consequent, likely dimerization pathway for diradical **3**.



Given the strong delocalization of the phenoxy spin density on the stilbenic olefin units, system **3** is probably more like a *meta*-benzoquinodimethane than a bis-phenoxy (Scheme 2). The position of highest spin density in **3** is not sterically protected, hence the diradical is so reactive that it is not observed when generated in solution. Apropos, one must parenthetically note that organic polyradical synthesis for magnetic materials involves a delicate balance between spin delocalization that is desirable because it promotes stronger exchange interaction between spin sites, and excessive spin delocalization that leads to undesirable radical couplings and formation of defects in the polyradical.

Assuming that **3** dimerizes as shown in Scheme 2, the resulting product should show hfc very much like those described for **20** (R=*tert*-Bu), but lacking one of the 0.5-0.6 G hyperfine couplings. This is exactly what is observed. Based on this analysis, we assigned the hfc as shown in Scheme 3 for **20**(R=*tert*-Bu) and the putative dimer of biradical **3**, structure **21**. The ability of **21** to survive further reaction to completely ESR-silent products is consistent with the similar persistence of **20** (R=*tert*-Bu) in solution. Since we do not see any evidence of biradical exchange in the spectrum from the oxidation of **19**, we presume that interruption of the conjugation pathway precludes exchange if two radical sites in **21** are simultaneously present.

Scheme 3. Hfc assignments for model systems and **21**. Hfc computed by UB3LYP/EPR-II method for **20**(H) are red. See [10] and [16] for literature descriptions of ESR for **20** (*t*-Bu). See Scheme 2 above for full, proposed structure of **21**.



Conclusions

We have synthesized several conjugated phenoxy-based conjugated molecules having direct conjugation from the phenoxy ring through the *para*-position to a second phenyl ring, with further linkage at the meta position. ESR hyperfine analysis and spectral analysis of related biradical model **1** shows that unpaired spin density delocalization does not occur to a significant extent past the phenyl ring into the *meta*-linked substituents. This does not rule out possible use of such building blocks for polyradical synthesis, but shows that exchange between such units would not be strong. Use of a *para*-styryl substituted phenoxy ring led to very strong delocalization with major spin density present on the ethenyl unit of the styryl group, as confirmed by computation and by analysis of model compounds. Diradical **3** based on the *para*-styryl conjugated unit was too unstable to observe – instead, we observed spectra consistent with a putative dimer that contains a persistent *para*-styryl phenoxy group. This second strategy appears to give too much spin delocalization to be useful for conjugated polyradicals, at least without significant modification to improve the stability of the strongly spin-bearing styryl segment of the radical units.

Acknowledgements.

This work was supported by the National Science Foundation (CHE 9521954 and CHE 9809548). Mass spectral analyses and elemental microanalyses were carried out at the University of Massachusetts Amherst Mass Spectrometry Facility and Microanalysis Facility, respectively.

Experimental

General

All $^1\text{H-NMR}$ spectra were obtained using either an IBM Instruments NR-80A (80 MHz) or a Bruker AC-200 (200 MHz) instrument. All spectra were recorded using chloroform- d_6 or acetone- d_6 and were calibrated using tetramethylsilane as internal standard. All IR spectra were recorded on a Perkin Elmer PE-1420 spectrometer or a Midac M-2000 FTIR with computer interface. UV-Vis spectra were recorded on a Shimadzu UV-260 double beam spectrometer. Melting points were obtained on Electrothermal IA6404 melting point apparatus, and are reported without correction. ESR spectra were obtained on a Bruker ESP-300 X-band (~ 9 GHz) spectrometer with standard computer interface and cooled nitrogen-gas variable temperature attachment. Solutions of stable radicals for ESR determination were subjected to three cycles of freeze-vacuum-thaw in order to remove oxygen, using quartz 4 mm O.D. tubes with a stopcock on top. All the reactions were run under an atmosphere of argon unless otherwise stated. All chemicals were purchased from Aldrich Chemical Company except where noted. All solvents were purchased from Fisher Scientific except where noted and were used without further purification except where noted. Dry tetrahydrofuran (THF) was distilled freshly under argon from sodium-benzophenone after pre-drying over lithium aluminum hydride. Anhydrous diethyl ether was either used directly from a freshly opened container or was distilled from potassium or sodium-benzophenone. Dry methylene chloride was distilled from calcium chloride. Dry triethylamine was distilled from phosphorus pentoxide. Dry dimethylformamide (DMF) was either purchased from Aldrich Chemical Company in a SureSealTM bottle or freshly distilled from phosphorus pentoxide.

Dimethyl 5-iodoisophthalate.

5-Iodoisophthalic acid (0.7 g, 2.4 mmol) was placed in a round bottom flask together with methanol (20 mL) and conc sulfuric acid (0.5 mL), and the mixture was refluxed overnight. Most of the methanol was removed under reduced pressure and the residue was dissolved in ethyl acetate. The organic layer was washed with water until the wash was neutral and dried over magnesium sulfate. A brown solid obtained after removal of the organic solvent was found to be a mixture. Silica gel column chromatography with 20/80 ethyl acetate/hexane separated the desired compound as a white crystalline solid (0.5 g, 65 %), mp 103-104.5 °C (lit. [17] 100-102 °C); $^1\text{H-NMR}$ (200 MHz, CDCl_3) δ 8.64-8.63

(t, 1H, J=1.46 Hz), 8.56-8.55 (d, 2H, J=1.44 Hz), 3.95 (s, 6H); MS-EI (m/z): calc'd for C₁₀H₉IO₄: 320; found: 320.

5-Iodoisophthalaldehyde.

Into a suspension of lithium aluminum hydride (0.62 g, 16.3 mmol) in dry diethyl ether (20 mL) cooled in an ice bath was added via a syringe a solution of dimethyl 5-iodoisophthalate (1.74 g, 5.44 mmol) in diethyl ether (5 mL). After the addition, the reaction mixture was stirred at room temperature for two more hours. The mixture was then poured into ice and extracted with ethyl acetate. The organic layer was washed with brine and dried over magnesium sulfate. The product, *1,3-dihydroxymethyl-5-iodobenzene* was obtained as a yellowish powder (0.94 g, 65 %) after removal of the solvent, and was used without further purification for oxidative conversion to 5-isophthalaldehyde. ¹H-NMR (200 MHz, CDCl₃) δ 7.62 (m, 2H), 7.34 (m, 1H), 4.69 (s, 4H) [18]. A mixture of the dihydroxy product (0.8 g, 3.03 mmol) from the previous step was heated under reflux for 2 h with pyridinium chlorochromate (1.95 g, 9.09 mmol) in dichloromethane (20 mL). TLC showed disappearance of the starting material and presence of a major spot with higher R_f. The reaction mixture was adsorbed on silica gel and eluted with 20/80 ethyl acetate/hexane. The desired compound was obtained after removal of solvent as a white crystalline solid with mp 117-119 °C (0.6 g, 76 %); ¹H-NMR (200 MHz, CDCl₃) δ 10.03 (s, 2H), 8.46-8.45 (d, 2H, J=1.44 Hz), 8.34-8.32 (t, 1H, J=1.44 Hz); IR (neat, cm⁻¹): 1690.

3,5-Di-*tert*-butyl-4-trimethylsiloxyphenyl boronic acid (6).

n-Butyllithium (3.7 mL of a 1.6 M solution in hexane, 5.9 mmol) was added into dry THF (10 mL) and the resulting yellow solution was cooled to -78 °C in a Dry Ice-acetone bath. A solution of 4-bromo-2,6-di-*tert*-butyltrimethylsiloxyphenol [19] (2.0 g, 5.6 mmol) in dry THF (15 mL) was added in via a cannula and the mixture was stirred at -78 °C for 20 min. Triisopropylborate (1.14 g, 6.1 mmol) was added via a syringe and stirring was continued for twenty more minutes. The mixture was then warmed up to room temperature, poured into a solution of hydrochloric acid (2 mL hydrochloric acid in 100 mL water), stirred for 2 h and extracted with ethyl acetate. The organic layer was washed with water and dried over magnesium sulfate. A yellowish powder was obtained after removal of the organic solvent. Recrystallization from 60/40 ethyl acetate/hexane gave a white crystalline solid (1.33 g, 74 %) with mp 230 °C (decomp.) (lit. mp 235-237 °C for the cyclic boroxin trimer of **6** [20]). ¹H-NMR (200 MHz, CDCl₃) δ 8.16 (s, 2H), 1.49 (s, 18H), 0.45 (s, 9H).

3,5-Di-*tert*-butyl-3'-formyl-4-hydroxybiphenyl (7).

Compound **6** (1.53 g, 4.73 mmol), 3-iodobenzaldehyde (1.0 g, 4.3 mmol), palladium acetate (48 mg, 5 mol %), tri-*o*-tolylphosphine (0.13 g, 10 mol %) and potassium carbonate (2.97 g, 21.5 mmol) were placed in a 100-mL round bottom flask, which was evacuated for 30 min and refilled with argon.

Dry DMF (15 mL) was added via a syringe. The mixture was heated at 80 °C for 5 h. The resulting green mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with water and dried over magnesium sulfate. The residue after removal of the organic solvent was applied to a silica gel column and eluted with 20/80 ethyl acetate/hexane. The main fraction was separated, the solvent evaporated and the solid residue recrystallized from hexane. A white crystalline solid (0.473 g, 35 %) was obtained, mp 155-157 °C; ¹H-NMR (200 MHz, CDCl₃) δ 10.09 (s, 1H), 8.04-8.03 (t, 1H, J=1.44 Hz), 7.84-7.79 (dquart, 2H, J=7.94 Hz, 1.8 Hz), 7.62-7.54 (t, 1H, J=7.94 Hz), 7.41 (s, 2H), 5.35 (s, 1H) 1.51 (s 18H); IR (KBr, cm⁻¹): 3640, 1690; UV-Vis (benzene, λ_{max}, nm): 290.6, 316.8; Elemental analysis: calc'd for C₂₁H₂₆O₂: C 81.29, H 8.39; found: C 81.55, H 8.52.

3,5-Di-tert-butyl-4-hydroxy-3'-vinylbiphenyl (**8**).

Methyltriphenylphosphonium bromide (0.26 g, 0.71 mmol) was suspended in THF (15 mL) and cooled in an ice bath. *tert*-Butyllithium (0.46 mL of a 1.7 M solution in hexane, 0.78 mmol) was added via a syringe giving a bright yellow solution. A solution of **7** (0.1 g, 0.32 mmol) in THF (5 mL) was added dropwise via a pressure-equalizing addition funnel. The color of the reaction turned to brown after the addition. The mixture was extracted with ethyl acetate and the organic layer was washed with water and dried over magnesium sulfate. The residue after removal of the organic solvent was applied to a silica gel column and eluted with 20/80 ethyl acetate/hexane. The desired compound was obtained as a yellow oil that crystallized into the yellow solid **8** (0.08 g, 80 %), with mp 72-73.5 °C. ¹H-NMR (200 MHz, CDCl₃) δ 7.55-7.26 (m, 6H), 6.86-6.72 (dd, 1H, J=17.34 Hz, 10.82 Hz), 5.85-5.76 (dd, 1H, J=17.68 Hz, 1.08 Hz), 5.31-5.26 (dd, 1H, J=10.84 Hz, 0.74 Hz), 5.28 (s, 1H), 1.50 (s, 18H). IR (neat, cm⁻¹): 3636, 2957, 989, 906. UV-Vis (benzene, λ_{max}, nm): 279.0. Elemental analysis: calc'd for C₂₂H₂₈O: C 85.71, H 9.09; found: C 84.86, H 9.24.

3,5-Di-tert-butyl-3',5'-diformyl-4-hydroxybiphenyl (**9**).

Compound **6** (1.24 g, 3.85 mmol), 5-iodoisophthalaldehyde (1.0 g, 3.85 mmol), palladium acetate (43 mg, 5 mol-%), tri-*o*-tolylphosphine (0.12 g, 10 mol-%) and potassium carbonate (1.6 g, 11.6 mmol) were placed in a three-neck round bottom flask, which was evacuated for 30 min and refilled with argon. Dry DMF (15 mL) was added via a syringe and the resultant yellow mixture was heated at 80 °C for 30 min. Upon heating, the reaction color changed from yellow to dark brown. TLC showed disappearance of the starting materials. The mixture was next poured into water and extracted with ethyl acetate. The organic layer was washed with water and brine thoroughly and dried over magnesium sulfate. The residue after removal of the organic solvent was applied to a silica gel column and eluted with 20/80 ethyl acetate/hexane. The desired product was isolated as a white crystalline solid (0.55 g, 42 %), mp 159-161 °C; ¹H-NMR (200 MHz, CDCl₃) δ 10.17 (s, 2H), 8.29 (m, 3H), 7.44 (s, 2H), 5.43 (s, 1H), 1.52 (s, 18H); IR (neat, cm): 3632, 1699; UV-Vis (benzene, λ_{max}, nm): 292; Elemental analysis: calc'd for C₂₂H₂₆O₃: C 78.11, H 7.69; found: C 78.23, H 7.74.

3,5-Di-tert-butyl-3',5'-divinyl-4-hydroxybiphenyl (10).

Into a suspension of methyltriphenylphosphonium bromide (0.35 g, 0.98 mmol) in THF (15 mL) cooled in an ice bath was added *tert*-butyllithium (0.63 mL of a 1.7 M solution in hexane, 1.07 mmol). A bright yellow color formed. Compound **9** (0.1 g, 2.96 mmol) in THF (2 mL) was added slowly via a syringe giving a pale yellow color. The mixture was stirred at room temperature for two more hours. TLC showed the main product was blue under short wave ultraviolet light and had a high R_f . The mixture was then poured into water, acidified with hydrochloric acid and extracted with ethyl acetate. The organic layer was washed with water, dried over magnesium sulfate, and evaporated. The residue was chromatographed (silica gel, 20/80 ethyl acetate /hexane) to give the product as a yellow crystalline solid (0.80 g, 81 %), mp 92-94 °C; $^1\text{H-NMR}$ (200 MHz, CDCl_3) δ 7.44-7.38 (m, 5H), 6.86-6.72 (dd, 2H, $J=17.32$ Hz, 10.82 hH), 5.87-5.78 (dd, 2H, $J=17.68$ Hz), 5.33-5.27 (dd, 2H, $J=0.72$ Hz), 5.29 (s, 1H), 1.50 (s, 18H); IR (neat, cm^{-1}): 3631, 2957, 988, 907; UV-Vis (benzene, λ_{max} , nm): 278.0; Elemental analysis: calc'd for $\text{C}_{24}\text{H}_{30}\text{O}$: C 86.22, H 8.98; found: C 85.99, H 9.00.

3,5-Di-tert-butyl-4-hydroxybiphenyl-3',5'-dicarboxylic acid (11).

Compound **6** (0.13 g, 0.403 mmol), 5-iodoisophthalic acid (0.11 g, 0.38 mmol), palladium acetate (10 mg, 5 mol-%) and potassium carbonate (1.0 g, 7.25 mmol) were added to a three-neck round bottom flask that was flushed with argon. A 1:1 mixture of acetone-water (40 mL) was added by a syringe and the mixture was heated at reflux under argon for 2 h. A black green mixture resulted. Acetone was removed under reduced pressure and the residue was extracted three times with ethyl ether and the organic layer was discarded. The aqueous layer was acidified with hydrochloric acid and then extracted with ethyl ether. The ether layer was washed with water and dried over magnesium sulfate and a yellowish solid (70 mg, 50%) was obtained after removal of solvent, mp 290 °C (d); $^1\text{H-NMR}$ (200 MHz, acetone- d_6) δ 8.61-8.60 (t, 1H, $J=1.6$ Hz), 8.43-8.42 (d, 2H, $J=1.6$ Hz), 7.52 (s, 2H), 1.52 (s, 18H); IR (KBr, cm^{-1}): 3630, 3449, 2922, 1702; UV-Vis (benzene, λ_{max} , nm): 290. A small amount of the sample was dissolved in acetone and the solvent was allowed to evaporate slowly to yield single crystals. X-ray diffraction analysis carried out by Prof. Clifford George of the Laboratory for the Structure of Matter at the Naval Research Laboratory. The structure was found to be the product in 1:1 ratio with acetone [14]. Elemental analysis: calc'd for $\text{C}_{22}\text{H}_{26}\text{O}_5 \cdot \text{C}_3\text{H}_6\text{O}$: C 70.09, H 7.48; found: C 70.00, H 7.59. Details of the crystal structure are available through the Cambridge Crystallographic Databank (CCDC Deposition # 137194).

Dimethyl 3,5-di-tert-butyl-4-hydroxybiphenyl-3',5'-dicarboxylate (12).

3,5-Di-*tert*-butyl-4-trimethylsiloxyphenyl boronic acid (**6**, 0.51 g, 1.56 mmol), dimethyl 5-iodoisophthalate (0.5 g, 1.56 mmol), palladium acetate (18 mg, 5 mol %), tri-*o*-tolylphosphine (95 mg, 20 mol %) and potassium carbonate (1.1 g, 8.0 mmol) were added to a three-neck round bottom

flask that was flushed with argon for 20 min. Dry DMF (30 mL) was added, and the mixture was heated at 70 °C for 1 h. A dark-brown color resulted. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed thoroughly with water and brine to remove DMF and dried over magnesium sulfate. The residue after removal of solvent was applied to a silica gel column and the main product was isolated as a yellowish solid (0.46 g, 77%), mp 160-162 °C. ¹H-NMR(200 MHz, acetone-*d*₆) δ 8.61-8.59 (t, 1H, J=2.0 Hz), 8.38-8.37 (d, 2H, J=2.0 Hz), 7.41 (s, 2H), 5.36 (s, 1H), 3.98 (s, 6H), 1.51 (s, 18H). UV-Vis (benzene, λ_{max}, nm): 284.2. Elemental analysis: calc'd for C₂₄H₃₀O₅: C 72.36, H 7.54; found: C 72.60, H 7.69.

Oxidation of 7.

A small amount of **7** (ca. 5 mg) was dissolved in benzene (1 mL) and lead dioxide was added. A dark green solution resulted, which was used for ESR and UV-Vis determination. UV-Vis (benzene, λ_{max}, nm): 350, 482, 599; ESR (9.651 GHz, benzene, gauss): a(H) = 1.80 (2H), 1.60 (2H), 1.86, 0.68.

Oxidation of 8.

A small amount of **8** (ca. 5 mg) was dissolved in benzene (1 mL) under argon and lead dioxide powder. The solution color changed to red-purple immediately and this solution was used for spectral analysis. UV-Vis (benzene, λ_{max}, nm): 328, 355, 507; ESR (9.670 GHz, benzene, gauss): a(H) = 1.76 (2H), 1.70 (2H), 1.73, 0.64.

Oxidation of 9.

A small amount of **9** (ca. 5 mg) was dissolved in benzene (1 mL) and excess lead dioxide powder was added. A dark green solution resulted, which was used for ESR and UV-Vis determination. UV-Vis (benzene, λ_{max}, nm): 341, 458, 615; ESR (9.652 GHz, benzene, gauss): a(H) = 1.77 (2H), 1.65 (2H), 1.78.

Oxidation of 10.

A small amount of **10** (ca. 5 mg) was dissolved in benzene (1 mL) under argon and excess lead dioxide powder was added. A red purple solution was formed at once. The solution was used directly for ESR. UV-Vis (benzene, λ_{max}, nm): 276, 337, 514; ESR (9.664 GHz, benzene, gauss): a(H) = 1.68 (2H), 1.64 (2H), 1.92.

Oxidation of 11.

To a dilute benzene solution of the radical precursor **11**, prepared by gentle warming and recooling, was added lead dioxide powder. The solution color changed to greenish yellow. The solution was used at once for spectral investigation. UV-Vis (benzene, λ_{\max} , nm): 267, 358, 505; ESR (9.795 GHz, benzene, gauss): a(H) = 1.83 G (2H), 1.72 G (2H), 1.62.

Oxidation of 12.

A small amount of **12** (ca. 5 mg) was dissolved in benzene (1 mL) and lead dioxide powder was added. A blue color formed immediately. The resulting blue solution was used for ESR and UV-Vis determinations. UV-Vis (benzene, λ_{\max} , nm): 347, 475, 615; ESR (9.800 GHz, benzene, gauss): a(H) = 1.70 (2H), 1.69 (2H), 1.69.

4-Acetoxy-3,5-di-tert-butyl-4'-methoxybiphenyl (13).

4-Bromo-2,6-di-tert-butylphenyl acetate [21] (2.0 g, 6 mmol), 4-methoxyphenyl boronic acid (**8**) (0.93 g, 6 mmol), palladium acetate (68 mg, 5 mol-%), tri-*o*-tolyl-phosphine (280 mg, 10 mol %) and potassium carbonate (1.28 g, 9.2 mmol) were added in a three-neck round bottom flask that was flushed with argon. Dry DMF (20 mL) was injected via a syringe and a greenish mixture formed. The mixture was heated at 85 °C for 2 h and a brown mixture afforded. The mixture was cooled to room temperature and poured into water (100 mL). The resulting mixture was extracted with ethyl acetate. The organic layer was washed thoroughly with water and brine to remove DMF and dried over magnesium sulfate. TLC showed the main product as a blue spot under short wavelength ultraviolet light. The red oil obtained after removal of solvent under reduced pressure was applied to a silica gel column and eluted with 50/50 hexane/methylene chloride. The major product was separated out as a yellow solid (1.38 g, 64%) after removal of solvent under reduced pressure. mp 87-90 °C; ¹H-NMR (200 MHz, CDCl₃) δ 7.50-7.46 (m, 4H), 6.99-6.95 (dd, 2H, J=8.0 Hz, 2.0 Hz), 3.85 (s, 3H), 2.37 (s, 3H), 1.39 (s, 18H).

4-Acetoxy-3'-bromo-3,5-di-tert-butyl-4'-methoxybiphenyl (14).

Compound **13** (0.24 g, 0.68 mmol) was dissolved in dichloromethane (10 mL), and Br₂ (0.33 g, 2 mmol) in dichloromethane (10 mL) was added. The red-brown solution was refluxed under argon overnight. Saturated sodium bisulfite solution was added to quench excess bromine and the organic layer was washed with water and brine several times. The crude solid obtained after removal of solvent was recrystallized from pentane to give prism-shaped white crystals (0.17 g, 57%) with mp 141-143 °C; ¹H-NMR (200 MHz, CDCl₃) δ 7.73-7.72 (d, 1H, J=2.0 Hz), 7.48-7.43 (dd, s, 3H, J=2.0 Hz), 6.98-

6.94 (d, 1H, J=8.0 Hz), 3.94 (s, 3H), 2.38 (s, 3H), 1.39 (s, 18H); Elemental analysis: calc'd for C₂₃H₂₉O₃Br: C 63.74, H 6.70; found: C 63.73, H 6.69.

1,3-Bis(4-Acetoxy-3,5-di-tert-butyl-4'-methoxybiphenyl-3'-(E)-β-ethenyl)benzene (15).

Compound **14** (0.5 g, 1.15 mmol), palladium acetate (7.8 mg, 3 mol %) and tri-*o*-tolylphosphine (21 mg, 6 mol %) were placed in a 50-mL round bottom flask. The flask was evacuated for 10 min and refilled with argon. A solution of tributylamine (0.24 g, 1.3 mmol) and 1,3-divinylbenzene (0.075 g, 0.58 mmol) in DMF (20 mL) was then added via syringe. The resulting yellow mixture was heated to ca 110 °C under argon overnight and a dark green color was found. TLC showed disappearance of starting materials. The mixture was poured into water and extracted with ethyl acetate. The organic layer was washed thoroughly with water and brine several times and dried over magnesium sulfate. After evaporating the solvent, the residue was applied onto a silica gel column and eluted with 20/80 ethyl acetate/hexane. The material with highest R_f was isolated after removal of solvent to give a white crystalline solid (0.21 g, 44 %) that was suitable for use in the subsequent deprotection step to make **16**. ¹H-NMR (200 MHz, CDCl₃) δ 7.76-6.94 (m, 18H), 3.95 (s, 6H), 2.39 (s, 6H), 1.41 (s, 36H); IR (KBr, cm⁻¹): 1760, 960; UV-Vis (THF, λ_{max}, nm): 214, 274, 3412.

1,3-Bis(3,5-Di-tert-butyl-4-hydroxy-4'-methoxybiphenyl-3'-(E)-ethenyl)benzene (16).

To a suspension of lithium aluminum hydride (0.1g, 2.6 mmol) in 10 mL of dry THF cooled in an ice bath, a solution of compound **15** (0.21 g, 0.25 mmol) in THF (5 mL) was added slowly via a cannula. A green suspension resulted and the reaction was allowed to stir for three hours at room temperature. The reaction was quenched by pouring into crushed ice and then acidified with hydrochloric acid. The resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, then dried over magnesium sulfate. An oily material was obtained after removal of solvent. Silica gel column chromatography with 20/80 ethyl acetate/hexane afforded a slightly greenish crystalline solid (0.1 g, 53 %) which sinters upon heating, mp 160 °C (dec). ¹H-NMR (200 MHz, CDCl₃) δ 7.73-6.95 (m, 18H), 5.25 (s, 2H), 3.94 (s, 6H), 1.51 (s, 36H). IR (neat, cm⁻¹): 3600, 960. UV-Vis (benzene, λ_{max}, nm): 284, 344. HRMS (EI, m/z): calc'd for C₅₂H₆₂O₄: 750.4648; found: 750.4651.

Oxidation of compound 16.

A small amount of compound **6** (ca. 5 mg) was dissolved in benzene (1 mL) under argon. Lead dioxide powder was added into this benzene solution and mixture was allowed to sit for a few minutes. A deep blue color was formed almost immediately. The final blue solution was separated from black lead dioxide by filtration. UV-Vis (benzene, λ_{max}, nm): 319, 372, 614; ESR (9.8 GHz, benzene, gauss): a(H) = 1.36 (2H), 1.92 (2H).

2,6-Di-tert-butyl-4-vinylphenyl acetate (17).

To a suspension of $\text{CH}_3\text{PPh}_3\text{Br}$ (15.25 g, 42.7 mmol) in THF (200 mL) cooled in an ice bath was added *n*-butyllithium (27.5 mL of a 1.6 M in hexane, 44.0 mmol) via a syringe and a yellow mixture resulted. The mixture was warmed up to room temperature and stirred for additional 30 min. A solution of 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde (5.0 g, 21.4 mmol) in THF was added dropwise via a pressure-equalizing addition funnel. The resulted mixture was stirred for 2 h. The color of the mixture changed to blue after two hours. The mixture was recooled in an ice bath and acetyl chloride (3.35 g, 42.7 mmol) was injected into the reaction mixture with a syringe. The reaction was warmed up to room temperature. The resulting brown mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with water and dried over magnesium sulfate. Silica gel column chromatography (20:80 ethyl acetate/hexane) separated out the desired product as an orange crystalline solid (3.53 g, 60 %) with mp 59-61 °C (lit mp 67 °C [22]); $^1\text{H-NMR}$ (200 MHz, CDCl_3) δ 7.36 (s, 2H), 6.76-6.62 (dd, 1H, $J=10.8$ Hz, 17.3 Hz), 5.72-5.62 (dd, 1H, $J=0.72$ Hz, 17.3 Hz), 5.24-5.18 (dd, 1H, $J=1.1$ Hz, 10.8 Hz), 2.35 (s, 3H), 1.36 (s, 18H); IR (KBr, cm^{-1}): 1760, 998, 905.

1,3-Bis[2-(3,5-di-tert-butyl-4-acetyloxyphenyl)ethenyl]benzene (18).

Compound **17** (0.2 g, 0.73 mmol), 1,3-diiodobenzene (0.12 g, 0.37 mmol), palladium acetate (8.2 mg, 5 mol %) and tri-*o*-tolylphosphine (22.2 mg, 10 mol %) were placed into a three-neck round bottom flask, which was evacuated for 30 min and refilled with argon. Triethylamine (1 mL) and DMF (10 mL) were added via syringe. The yellow mixture was heated under argon at 100 °C. After overnight reaction, a dark colored mixture was produced. TLC showed that the starting materials were gone and that the main product spot fluoresced blue under short wavelength UV light. The reaction was quenched by pouring into water and the resulting mixture extracted with ethyl acetate. The organic layer was washed with dilute hydrochloric acid and then water several times to remove DMF. Careful separation by column chromatography (10:90 ethyl acetate/hexane) gave the desired product **18** as a yellowish solid (0.157 g, 70%) that was suitable for use in the subsequent deprotection step to make **19**. $^1\text{H-NMR}$ (200 MHz, CDCl_3) δ 7.65-7.33 (m, 18H), 7.20-7.12 (d, 2H, $J=16$ Hz), 7.07-6.99 (d, 2H, $J=16$ Hz), 2.36 (s, 6H), 1.39 (s, 36H); IR (neat, cm^{-1}): 2956, 2850, 1760, 960.

1,3-Bis[2-(3,5-di-tert-butyl-4-hydroxyphenyl)ethenyl]benzene (19).

To an ice cooled suspension of lithium aluminum hydride (0.055 g, 1.46 mmol) in dry THF (2 mL) was added dropwise a solution of compound **18** (0.157 g, 0.25 mmol) in dry THF (2 mL). The color of the suspension changed from grey to yellowish and it was allowed to stir at room temperature for an additional 4 h. The mixture was poured into ice, acidified with conc HCl and extracted with ethyl acetate. The organic layer was washed with water and dried over magnesium sulfate. A yellow oil was obtained after removal of the solvent. Silica gel column chromatography gave a fraction that contained

mainly the desired product with some impurities. Slow evaporation of the solvent allowed growth of yellow crystals (0.1 g, 74 %, mp 105 °C[d]) that showed only one spot by TLC. ¹H-NMR (200 MHz, CDCl₃) δ 7.63 (m, 1H), 7.37 (m, 7H), 7.18-7.10 (d, 2H, J=16), 7.00-6.92 (d, 2H, J=16), 5.30 (s, 2H), 1.49 (s, 36H); IR (neat, cm⁻¹): 3638, 2956, 2871, 957; UV-Vis (benzene, λ_{max}, nm): 328 nm; Elemental analysis: calc'd for C₃₈H₅₀O₂: C 84.76, H 9.29; found: C 85.20, H 9.69.

Oxidation of **19**.

A small amount of compound **19** (ca. 5 mg) was dissolved in dry benzene (1 mL) under argon. To this solution was added lead dioxide powder. Color changed from colorless to dark yellow. Lead dioxide was filtered off and the filtrate was subjected ESR spectroscopy. IR showed disappearance of the sharp OH stretch at 3640 cm⁻¹ that is observed in compound **19**. UV-Vis (benzene, λ_{max}, nm): 309, 457, 488; ESR (9.751 GHz, benzene, gauss): a(H) = 6.37, 2.93, 1.69, 1.59, 1.48, 1.47, 1.40, 0.53.

References and Notes

- (a) Mataga, N. "Possible 'Ferromagnetic States' of Some Hypothetical Hydrocarbons" *Theor. Chim. Acta* **1968**, *10*, 372-376; (b) Ovchinnikov, A. A. "Multiplicity of Ground-State of Large Alternant Organic Molecules with Conjugated Bonds" *Theor. Chim. Acta* **1978**, *47*, 297-304; (c) Borden, W. T.; Davidson, E. R. "Electron Repulsion in Diradicals" *J. Am. Chem. Soc.* **1977**, *99*, 4587; (d) Borden, W. T., Ed.; *Diradicals*; John Wiley and Sons, Inc.: New York, 1982.
- Rajca, A. "Organic Diradicals and Polyradicals: From Spin Coupling to Magnetism?" *Chem. Rev.* **1994**, *94*, 871.
- (a) Rajca, A.; Wongsriratanakul, J.; Rajca, S. "Magnetic ordering in an organic polymer" *Science* **2001**, *294*, 1503-1505; (b) Rajca, A. "From high-spin organic molecules to organic polymers with magnetic ordering" *Chem. Eur. J.* **2002**, *8*, 4834-4841.
- (a) Koga, N.; Inoue, K.; Sasagawa, N.; Iwamura, H. "Magnetic properties of microcrystalline "poly(phenyldiacetylenes)" carrying radical or carbene centers on the side chains" *Mater. Res. Soc. Symp. Proc.* **1990**, *173*, 39-49; (b) Abdelkader, M.; Drenth, W.; Meijer, E. W. "Synthesis and Characterization of a Stable Poly(iminomethylene) with Pendant Phenoxy Radicals" *Chem. Mater.* **1991**, *3*, 598; (c) Fujii, A.; Ishida, T.; Koga, N.; Iwamura, H. "Syntheses and magnetic properties of poly(phenylacetylenes) carrying a (1-oxido-3-oxy-4,4,5,5-tetramethyl-2-imidazolyl-2-yl) group at the meta or para position of the phenyl ring" *Macromolecules* **1991**, *24*, 1077-1082; (d) Miura, Y.; Inui, K.; Yamaguchi, F.; Inoue, M.; Teki, Y.; Takui, T.; Itoh, K. "Molecular design, synthesis, and magnetic characterization of poly(phenylacetylene) with pi-toporegulated pendant nitronyl nitroxide radicals as models for organic superpara- and ferromagnets" *J. Polym. Sci., Part A: Polym. Chem.* **1992**, *30*, 959-966; (e) Nishide, H. "High-spin alignment in pi-conjugated polyradicals. A magnetic polymer" *Adv. Mater.* **1995**, *7*, 937-941; (f) Nishide, H.; Ozawa, T.; Miyasaka, M.; Tsuchida, E. "A nanometer-sized high-spin polyradical: Poly(4-phenoxy-1,2-

- phenylenevinylene) planarily extended in a non-Kekulé fashion and its magnetic force microscopic images" *J. Am. Chem. Soc.* **2001**, *123*, 5942-5946; (g) Miyasaka, M.; Saito, Y.; Nishide, H. "Magnetic force microscopy images of a nanometer-sized purely organic high-spin polyradical" *Adv. Funct. Mater.* **2003**, *13*, 113-117.
- Lahti, P. M.; Ichimura, A. S. "A Semiempirical Study of Electron Exchange Interaction in Organic High-Spin Pi-Systems - Classifying Structural Effects in Organic Magnetic Molecules" *J. Org. Chem.* **1991**, *56*, 3030-3042.
 - Lahti, P. M. "The Design of Organic Based Materials with Controlled Magnetic Properties", In *Molecule-Based Magnetic Materials. Theory, Techniques, and Applications*; Turnbull, M. M.; Sugimoto, T.; Thompson, L. K., Eds.; American Chemical Society: Washington, DC, 1996; Vol. 644, pp. 218-235.
 - Nishide, H.; Miyasaka, M.; Tsuchida, E. "Average Octet Radical Polymer: A Stable Polyphenoxyl with Star-Shaped pi-Conjugation" *Angew. Chem. Int. Ed.* **1998**, *37*, 2400-2402.
 - (a) Modarelli, D. A.; Rossitto, F. C.; Lahti, P. M. "Convenient Unimolecular Sources of Aryloxyl Radicals. I. Aryloxyoxalyl Chlorides" *Tetrahedron Lett.* **1989**, *30*, 4473-4476; (b) Modarelli, D. A.; Rossitto, F. C.; Lahti, P. M. "Convenient Unimolecular Sources of Aryloxyl Radicals .2. Aryloxyoxalyl Tert-Butylperoxides" *Tetrahedron Lett.* **1989**, *30*, 4477-4480; (c) Modarelli, D. A.; Lahti, P. M. "Diaryl Oxalates as Convenient Unimolecular Sources of Aryloxyl Radicals" *Chem. Commun.* **1990**, 1167-1168; (d) Rossitto, F. C.; Lahti, P. M. "Poly[3,5-Di-Tert-Butyl-4-[(2,4,6-Tri-Tert-Butylphenyl)Oxalato]Phenylacetylene] – a Photochemical Precursor to a Conjugated Polyradical" *Macromolecules* **1993**, *26*, 6308-6309; (e) Lahti, P. M.; Modarelli, D. A.; Rossitto, F. C.; Inceli, A. L.; Ichimura, A. S.; Ivatury, S. "Aryl Oxalate Derivatives as Convenient Precursors for Generation of Aryloxyl Radicals" *J. Org. Chem.* **1996**, *61*, 1730-1738; (f) Lahti, P. M.; Inceli, A. L.; Rossitto, F. C. "Synthesis, Characterization, and Photolysis of Poly[3,5-tert-butyl-4-[(2,4,6-tri-tert-butylphenyl)oxalato]phenylacetylene], a Photochemical Polyradical Precursor" *J. Polym. Sci. Part A, Polym. Chem.* **1997**, *35*, 2167-2176.
 - (a) Wolf, S. A.; Awschalom, D. D.; Buhrman, R. A.; Daughton, J. M.; von Molnar, S.; Roukes, M. L.; Chtchelkanova, A. Y.; Treger, D. M. "Spintronics: A Spin-Based Electronics Vision for the Future", *Science* **2001**, 1488-1495; (b) Xiong, Z. H.; Wu, D.; Vardeny, Z. V.; Shi, J. "Giant magnetoresistance in organic spin-valves" *Nature* **2004**, *427*, 821 – 824.
 - Becker, H.-D. "New Stable Phenoxy Radicals. The Oxidation of Hydroxystilbenes." *J. Org. Chem.* **1969**, *34*, 1211-1215.
 - Duling, D. R. "Simulation of Multiple Isotropic Spin Trap EPR Spectra" *J. Magn. Res.* **1994**, *B104*, 105-110.
 - Weil, J. A.; Bolton, J. R.; Wertz, J. E. "Electron Paramagnetic Resonance: Elementary Theory and Practical Applications"; Wiley: New York, NY, 1986; pp. 182-186.
 - Cf. Goldman, J.; Petersen, T. E.; Torssell, K.; Becher, J. "¹⁹F and ¹H-NMR and ESR Investigations of Aryl-*t*-Butyl Nitroxides and Nitronylnitroxides" *Tetrahedron* **1973**, *29*, 3833-3842.

14. Lahti, P. M.; Esat, B.; Ferrer, J. R.; Liu, Y.; Marby, K. A.; Xie, C.; George, C.; Antorrena, G.; Palacio, F. "Polymeric, H-bonded, and chelatable phenoxyl and nitroxide radicals" *Mol. Cryst. Liq. Cryst.* **1999**, *334*, 285-294.
15. Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Robb, M. A.; Cheeseman, J. R.; Keith, T.; Petersson, G. A.; Montgomery, J. A.; Raghavachari, K.; Al-Laham, M. A.; Zakrzewski, V. G.; Ortiz, J. V.; Foresman, J. B.; Cioslowski, J.; Stefanov, B. B.; Nanayakkara, A.; Challacombe, M.; Peng, C. Y.; Ayala, P. Y.; Chen, W.; Wong, M. W.; Andres, J. L.; Replogle, E. S.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Binkley, J. S.; Defrees, D. J.; Baker, J.; Stewart, J. P.; Head-Gordon, M.; Gonzalez, C.; Pople, J. A. *Gaussian 98*; Gaussian Inc.: Pittsburgh, PA, 1998.
16. Brigati, G.; Lucarini, M.; Mugnaini, V.; Pedulli, G. F. "Determination of the Substituent Effect on the O-H Bond Dissociation Enthalpies of Phenolic Antioxidants by the EPR Radical Equilibration Technique" *J. Org. Chem.* **2002**, *67*, 4828-4832.
17. Sundberg, R. J.; Heintzelman, R. W. "Reactivity of aryl nitrenes. Competition between carbazole formation and internal bond reorganization in biphenylnitrenes" *J. Org. Chem.* **1974**, *39*, 2546-2552.
18. Zeng, F.; Zimmerman, S. C. "Rapid Synthesis of Dendrimers by an Orthogonal Coupling Strategy" *J. Amer. Chem. Soc.* **1996**, *118*, 5326-5327.
19. Harrer, W.; Kurreck, H.; Reusch, J.; Gierko, W. "Über galvinole und galvinoxyl-mehrspinsysteme—I : Eine neue metallorganische synthese von mono- und oligo-galvinolen" *Tetrahedron* **1975**, *31*, 625-632.
20. Satoh, Y.; Shi, C. "An Efficient Synthesis of 4-Aryl-2,6-di-*tert*-butylphenols by a Palladium Catalyzed Cross-Coupling Reaction" *Synthesis* **1994**, 1146-1148
21. Nekhoroshev, M. V.; Ivakhnenko, E. P.; Okhlobystin, O. Yu "Methods for the acylation of sterically hindered phenols" *Zh. Obshch. Khim.* **1977**, *13*, 662.
22. Braun, D.; Wittig, W. "Stable polyphenoxyls. 4. Polymers from sterically hindered, protected vinylphenols" *Makromol. Chem.* **1980**, *181*, 557-563.

Sample availability: Samples are not available.